




SST- 14 Regulation of External Constraints & The Emergence of Living Systems



On the early Earth, chemical reactions occur continuously.

Molecules collide, combine, and break apart. Energy flows through the environment, driving change. Yet most of these processes are short-lived. Reactions occur and then disperse, their products carried away or broken down.

Over time, some reactions begin to reinforce themselves. Certain combinations of molecules make it easier for similar reactions to occur again. These processes persist slightly longer, but they remain fragile, easily disrupted by changing conditions.

Then a further development occurs. Some of these reactions become enclosed within simple boundaries, small, membrane-like structures that hold their components together. What was once dispersed is now contained. Reactions occur within a protected space.

Within these boundaries, new processes emerge. Systems begin to take in materials from their surroundings and transform them in ways that sustain their own organisation. For the first time, persistence is no longer accidental. It becomes organised.

This marks the beginning of living systems.

Formal Description

External constraints arise from physical, energetic, and environmental conditions that define the requirements for system viability.

Early systems faced fundamental external constraints, including dispersion, energy dissipation, and environmental fluctuation, which limited their persistence.

The emergence of catalysis, autocatalysis, and autocatalytic sets reduced constraints on chemical processes, enabling more persistent organisation.



Compartmentalisation further altered constraint configurations by maintaining the proximity of interacting components and reducing environmental disruption. The emergence of metabolism introduced active regulation of external constraints through the acquisition and transformation of external satisfiers, giving rise to needs as an intrinsic feature of viability. These developments represent successive improvements in constraint regulation, enabling increasingly stable and persistent systems.

Plain English Explanation


At the most basic level, systems are constrained by the physical world: energy is lost; materials disperse; and conditions change. These factors make it difficult for organised systems to persist. Early chemical systems had no way of controlling these conditions. Reactions happened, but they did not last. For a system to persist, something had to change.

The first step was the emergence of processes that made certain reactions easier to sustain.

Catalysis allowed reactions to occur more readily. Autocatalysis allowed reaction networks to reinforce themselves. These developments made persistence more likely, but still fragile.

A major shift occurred with the emergence of boundaries. When processes became enclosed within structures such as simple membranes, the components of the system could remain together. This reduced the effects of environmental disruption and allowed processes to continue more reliably. The next major step was metabolism. Systems began to actively take in materials and energy from their surroundings and transform them in ways that sustained their organisation. At this point, persistence depended on ongoing access to specific inputs. Needs emerged because the system could no longer continue without them.

Living systems can therefore be understood as systems that actively regulate external constraints. They do not simply exist within their environment; they manage their relationship with it in order to persist.



Example 1 (Catalysis)

Certain chemical reactions occur more easily in the presence of catalysts. This reduces constraints on reaction rates and allows processes to continue more reliably.



Example 2 (Compartmentalisation)

A lipid vesicle forms a simple boundary that holds chemical components together, reducing dispersion and stabilising reactions.

Example 3 (Metabolism)

A living cell takes in nutrients and converts them into energy and structural components, actively maintaining its organisation.

Provenance and Links

This module draws on:

- Non-equilibrium thermodynamics, particularly Ilya Prigogine, which explains how organised systems persist under conditions of energy flow.
- The theory of major evolutionary transitions developed by John Maynard Smith and Eörs Szathmáry, which describes the emergence of new levels of biological organisation.
- Research on autocatalytic systems and the origins of life, including Stuart Kauffman.
- General systems theory, including Ludwig von Bertalanffy, which emphasises open systems and exchanges with the environment.

This module interprets these developments as successive improvements in the regulation of external constraints required for system viability.

Practical Exercise

Consider a simple living system (e.g. a plant cell or bacterium).

Explain:

1. What external constraints affect this system?
2. How does the system regulate these constraints?
3. What needs arise as a result of this regulation?

👉 Write a short paragraph (6–8 sentences).